

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

form of their material as were formerly their confreres of the Paleozoic, who needed nearly a hundred years to recognize that many of the arboreal forms of the carboniferous forests were not seed plants but vascular cryptogams, a view long urged by the anatomists and finally admitted on every hand. The flora of the Middle Eocene of Georgia is chiefly interesting because of the fact that it clearly marks, in the opinion of the author, a warmer climate for that epoch than for the early Eocene.—E. C. Jeffrey.

Phylogenetic taxonomy of angiosperms.—Professor Bessey for many years was interested in a phylogenetic scheme of classification, especially with reference to the angiosperms. In a series of papers he has developed his point of view, and his final paper was presented in connection with the twenty-fifth anniversary celebration of the Missouri Botanical Garden. 5 An interesting feature of the paper is the definite formulation of the principles of classification as applied to flowering plants. These principles are given in the form of 28 "dicta," 7 of them of general application, and the remainder having special reference to flowering plants. These dicta announce the primitive and derived condition in reference to numerous structures, some of them generally accepted, and some of them under discussion. The author then applies these numerous dicta to the taxonomic complexities of angiosperms. He substitutes "Oppositifoliae" and "Alternifoliae" for the old names dicotyledons and monocotyledons, on the theory that dicotyledons are primitively oppositeleaved, as shown by their cotyledons; and by the same sign monocotyledons are essentially alternate-leaved. With this start, and in this way, the whole sequence of angiosperms is presented in 300 families, beginning with Alismataceae and ending with Lactucaceae (the Compositae being presented as 14 families). It is a very laborious assembling of material on the basis of assumed phylogenetic sequences, quite comparable with Engler's Syllabus, and useful in the same way.—J. M. C.

Amoeboid movements of chromatophores.—The large epidermal leucoplasts of mature leaves of *Orchis latifolius* and *O. incarnatus*, according to Küster, ¹⁶ display amoeboid changes of form, having the power to put out and take in processes resembling pseudopodia. Also these leucoplasts at times break up into a large and small portion, which in turn may again fuse. Küster does not accept the idea of Senn that there is a peristromium surrounding the chromatophore, holding rather that the pseudopodia belong to the chromatophore itself. Sometimes the chromatophore as a whole moves in the direction taken by a pseudopodium, thus exhibiting active movement. More commonly, however, a leucoplast suffers no change in position, following a pseudopodial

¹⁵ Bessey, Charles E., The phylogenetic taxonomy of flowering plants. Annals Mo. Bot. Gard. 2:109-164. fig. 1. 1915.

¹⁶ KÜSTER, ERNST, Über amöboide Formveränderungen der Chromatophoren höherer Pflanzen. Ber. Deutsch. Bot. Gesells. **29**:362–369. figs. 4. 1911.